

BANOL®

TURF AND ORNAMENTAL FUNGICIDE For control of Pythium and Phytophthora

**Only for Use on Golf Courses, Sod Farms, Commercially Grown Ornamentals in Greenhouses
and Container-Grown Ornamentals in Nurseries**

NOT FOR USE ON SOD FARMS IN ARIZONA

ACTIVE INGREDIENT:

Propamocarb hydrochloride*	66.5%
INERT INGREDIENTS:	<u>33.5%</u>
TOTAL	100.0%

*Propyl [3-(dimethylamino)propyl] carbamate monohydrochloride
Contains 6.0 lbs. active ingredient per gallon

EPA Reg. No. 432-942
EPA Est. No. 407-IA-02

NET CONTENTS:

KEEP OUT OF REACH OF CHILDREN **CAUTION** FIRST AID

<p>IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told by a poison control center or doctor. Do not give anything by mouth to an unconscious person.</p>
<p>IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing, eye. Call a poison control center or doctor for treatment advice.</p>
<p>IF ON SKIN or CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.</p>
<p>NOTE TO PHYSICIAN: Do not administer atropine unless cholinesterase inhibition is measured or patient shows distinct symptoms thereof. Oxime therapy is contraindicated. Otherwise, administration of gastric lavage or induction of vomiting followed by application of activated charcoal is the recommended treatment.</p>

<p>IN CASE OF MEDICAL EMERGENCIES OR HEALTH AND SAFETY INQUIRIES, OR IN CASE OF FIRE, LEAKING, OR DAMAGED CONTAINERS, INFORMATION MAY BE OBTAINED BY CALLING 1-800-228-5635, EXT. 202. HAVE THE PRODUCT CONTAINER OR LABEL WITH YOU WHEN CALLING A POISON CONTROL CENTER OR DOCTOR, OR GOING FOR TREATMENT.</p>
--

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers (both WPS and Non-WPS) must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove protective clothing and equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

USE PRECAUTIONS

Do not graze or feed clippings from treated areas to livestock.

Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container and keep tightly closed. Store in a cool dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DO NOT REUSE ORIGINAL CONTAINER.

GENERAL INFORMATION

BANOL[®] is a completely water miscible fungicide concentrate. BANOL is intended for use on turfgrass or ornamental species which are subject to Pythium Blight (Cottony Blight, Grease Spot) or *Phytophthora*. It may be used after germination to control Pythium damping-off in turf, or on established turf to prevent or control Pythium Blight. BANOL does not afford broad spectrum disease control for *Rhizoctonia* sp. (brown patch), *Sclerotinia* sp. (dollarspot), or *Helminthosporium* sp. (leafspot). Use a broad spectrum fungicide if these diseases coexist with Pythium Blight.

BANOL is also recommended for prevention of root rot and damping-off caused by *Pythium* sp. and *Phytophthora* sp. at all stages of ornamental propagation.

COMPATIBILITY: BANOL is compatible with benomyl, iprodione, ethazol, chloroneb, trichlorfon, triadimefon, chlorothalonil, diazinon, chlorpyrifos, and thiram.

For control of other soil borne diseases such as *Rhizoctonia* sp. or *Fusarium* sp. BANOL can be tank-mixed with benomyl or thiophanate-methyl. Use the full rate of each fungicide and follow the most restrictive directions and restrictions appearing on the labels of these products.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers, other persons, or pets, either directly or through drift. Keep people and pets out of the area during application. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants.
- Chemical-resistant footwear plus socks.
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant headgear for overhead exposure

Notify workers of the application by warning them orally and by posting signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter treated areas until sprays have dried.

TURFGRASS	AREA	Fluid Ounces BANOL	REMARKS
For Control of: Pythium Blight, Pythium Damping Off, and Root Rot on Established Turf and Overseeded Areas	1000 ft ²	<p>Preventative Treatment: 1¹/₃-2 fluid oz. in 2-5 gallons of water</p> <p>Curative Treatment: 3-4 fluid oz. in 2-5 gallons of water</p>	<p>Do not apply more than a total of 12 fl. oz. (0.57 lb a.i.) of BANOL per 1000 ft² of turfgrass per year, equivalent to 25 lb a.i./acre/year.</p> <p>ESTABLISHED TURF: Apply as a preventative treatment during periods of high temperature and humidity favorable to the development of Pythium Blight.</p> <p>OVERSEEDDED AREAS: Apply after germination to control Pythium Damping-Off.</p> <p>RETREATMENT: Re-treat at 7-21 day intervals if conditions remain favorable for disease development. Use lower rate for the shorter interval and the higher rate for the longer interval within the application rate range. Under severe disease pressure conditions, use the highest recommended rate and shortest interval.</p>

BANOL is recommended for prevention of root rot and damping-off caused by *Pythium* sp. and *Phytophthora* sp. at all stages of turfgrass development and ornamental plant propagation, including seeding, transplanting and potting.

ORNAMENTALS	Fluid Ounces BANOL	REMARKS
For Control of: <i>Pythium</i> and <i>Phytophthora</i> on Ornamental Plants: African Violet, Ageratum, Alyssum, Aster, Azalea, Browallia, Carnation, Celosia, Chrysanthemum, Coleus, Dahlia, Dusty Miller, Easter Lily, Ferns, Geranium, Gloxinia, Impatiens, Lobelia, Marigold, Pansy, Petunia, Poinsettia, Portulaca, Salvia, Snapdragon, Verbena, Violet, and Vinca.	3 fl. oz. per 10 gallons of water 3 fl. oz. per 10 gallons of water 2 fl. oz. per 10 gallons of water 2 fl. oz. per 10gallons of water	<p>Do not use for field grown ornamentals.</p> <p>BANOL requires no agitation after initial mixing. Stock solutions of BANOL should be used within one week of mixing. BANOL does not cause phytotoxic symptoms on the listed plants when used as directed. Trials on other ornamental species should be conducted before extensive use.</p> <p>Seeding: Apply to the seeding area using 3 quarts of solution/10 ft². Treat before sowing of seeds requiring light and after covering of seeds requiring dark for germination.</p> <p>Transplanting (Seedlings): Apply to the transplanting area using 3 quarts of solution /10 ft² before transplanting.</p> <p>Transplanting (Cuttings): Drench the cutting bed using 2 gallons of solution /10 ft² before setting.</p> <p>Potting: Dip bare root plants in the solution before potting. After potting drench each pot using 3.5 fl. oz. of solution per 4 inch pot. Repeat the drench treatment at 6 week intervals.</p>
Woody Ornamentals	2.5 fl. oz. per 10 gallons of water	For control of <i>Phytophthora sp.</i> on woody ornamentals, drench each container at the rate of 5 quarts of solution /10 ft ² of surface. Repeat the drench treatment at 3 to 4 month intervals

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of extreme weather and soil conditions, manner of use and other factors beyond Bayer Environmental Science control, it is impossible for Bayer Environmental Science to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability or otherwise.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE ELECTION, THE REPLACEMENT OF PRODUCT.



Bayer Environmental Science

Bayer Environmental Science
A Business Group of Bayer CropScience LP
PO Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, NC 27709