

TURF-SPECIFIC USE DIRECTIONS* (Continued)

Disease Control	Rate # of Product /1,000 sq ft	Use Information
Necrotic Ring Spot (Diplodia/Phoma)	0.38 - 0.50	Begin fungicide applications preventively when soil temperatures reach 60° F or early spring or summer and continue at 28 day intervals when conditions are favorable for development. Apply at sunrise or shortly after movement of fungicide into the root zone.
Pink Patch (Lophomyces coenophialii)	0.55 - 1.1	Begin fungicide applications preventively when soil temperatures reach 60° F to 70° F during periods of high rainfall when conditions are favorable for development. Apply preventively applications when light disease pressure is anticipated, use the lower rate and longer interval. Under conditions for severe disease or for an early curative application, use the higher rate and shorter interval.
Red Thread (Laetisaria fraxinea)	0.41 - 1.1	Begin fungicide applications preventively when night air temperatures reach 60° F to 70° F under periods of high rainfall when conditions are favorable for development. Under preventive applications when light disease pressure is anticipated, use the lower rate and longer interval. Under conditions for severe disease or for an early curative application, use the higher rate and shorter interval. Continue to monitor on a 14-28 day interval.
Rust (Puccinia spp.)	0.28 - 0.55	Begin fungicide applications preventively when conditions are favorable for development. Under preventive applications when light disease pressure is anticipated, use the lower rate and longer interval. Under conditions for severe disease or for an early curative application, use the higher rate and shorter interval. Continue to monitor on a 14-28 day interval.
Summer Patch (Magnaporthe poae)	0.55 - 1.1	Begin fungicide applications preventively when soil temperatures reach 60° F to 80° F at a 2-4 inch soil depth when warm and/or high humidity conditions are present. Continue on an interval, use the higher rate and shorter interval under severe disease pressure or if there is prior history.
Take-all Patch (Gaeumannomyces graminis var. zeae)	0.55 - 1.1	Begin fungicide applications preventively in the fall. Make 1-2 applications, depending on 14-28 day intervals when light disease conditions are favorable for development. Under preventive applications when light disease pressure is anticipated, use the lower rate and longer interval. Under conditions or prior history for severe disease or for an early curative application, use the higher rate and shorter interval.

7

Continued

TURF-SPECIFIC USE DIRECTIONS* (Continued)

Disease Control	Rate # of Product /1,000 sq ft	Use Information
Large Patch (Rhizoctonia Large Patch/Zeax/Phaen) (Rhizoctonia solani)	2.55 - 1.1	Begin fungicide applications preventively in the fall. Make 1-2 applications, depending on 14-28 day intervals when conditions are favorable for development. Under preventive applications when light disease pressure is anticipated, use the lower rate and longer interval. Under conditions or prior history for severe disease or for an early curative application, use the higher rate and shorter interval.
Additional Turfgrass Uses	Rate out/1,000 sq ft	Use Information
Algae	0.55 - 1.1	Use preventively the lower rate to the suppression of algae. Continue as needed, on a 14-30 day interval. Tank mix with chlorinated or non-chlorinated to improve the overall control of algae.
Summer stress syndrome / summer patch	0.55 - 1.1	Apply applications prior to the onset of conditions for abiotic stress to reduce symptoms of summer stress/summer patch. Chloro-Dimethane to improve the relief in symptoms on hot and/or moisture to severe stress.

DO NOT apply more than 3.3 fl oz per 1,000 sq ft of Chipco TRITON FLO per year.

PRODUCT QUANTITY (FL OZ) BY SPRAY VOLUME AND TANK CAPACITY
Chipco TRITON FLO at 0.55 fluid ounces per 1,000 sq ft

Spray Tank Capacity	spray volume (gallons per 1,000 sq ft)				
	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
25 Gallons	14.6 fl oz	7.0 fl oz	4.7 fl oz	3.5 fl oz	2.8 fl oz
50 Gallons	29.0 fl oz	14.0 fl oz	9.3 fl oz	7.0 fl oz	5.6 fl oz
100 Gallons	58.0 fl oz	28.0 fl oz	18.7 fl oz	14.0 fl oz	11.2 fl oz
200 Gallons	112.0 fl oz	56.0 fl oz	37.3 fl oz	28.0 fl oz	22.4 fl oz

8

Chipco TRITON FLO at 1.1 fluid ounces per 1,000 sq ft

Spray Tank Capacity	SPRAY VOLUME (gallons per 1,000 sq ft)				
	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
25 Gallons	29.0 fl oz	14.0 fl oz	9.3 fl oz	7.0 fl oz	5.6 fl oz
50 Gallons	58.0 fl oz	28.0 fl oz	18.7 fl oz	14.0 fl oz	11.2 fl oz
100 Gallons	112.0 fl oz	56.0 fl oz	37.3 fl oz	28.0 fl oz	22.4 fl oz
200 Gallons	224.0 fl oz	112.0 fl oz	74.6 fl oz	56.0 fl oz	44.8 fl oz

IMPORTANT: READ BEFORE USE
Read the entire Directions for Use, Conditions, Disclaimers of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimers of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, malfeasance or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP or its subsidiaries. Such risks shall be assumed by the user of Bayer.

DISCLAIMERS OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE. HAVE EXTENSIVE TESTING OF THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, DAMAGES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID BY OR BUYER CROPSCIENCE LP'S ELECTRONIC REPLACEMENT OF PRODUCT.

Bayer and G, the Bayer Cross and G, Chipco®, Chipco TRITON®, Compass®, 26 G®, Signature™, and Backed by Bayer™ are trademarks of Bayer.

9

PULL HERE TO OPEN



For control of diseases on turfgrasses of commercial, institutional and residential turfs, such as golf courses, and farms, commercial lawns, rights of way, parks, recreation areas and sports fields.

ACTIVE INGREDIENTS: Tritonazole..... 26.1%
OTHER INGREDIENTS:..... 73.9%
Total:..... 100.0%
 *Contains 3 lbs trifluralin per US gallon
 EPA Reg. No. 432-1487

KEEP OUT OF REACH OF CHILDREN
CAUTION

For MEDICAL and TRANSPORTATION Emergencies ONLY, call 24 Hours a Day 1-800-334-7377
 For PRODUCT USE Information Call 1-800-331-2697

Net Contents
1 Gallon
 79450140
 11/2005A

BACKED BY BAYER



Produced for:
Bayer Environmental Science
 A Business Group of Bayer CropScience LP
 2 T. W. Alexander Drive
 Research Triangle Park, NC 27709

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Never perspire a glass of water if able to swallow.
- Do not get anything in mouth by an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES OR CONTACT WITH EYES:

- Hold eyes open and flush with clean water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: Treat Symptomatically.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Hazard if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

Wear long-sleeved shirt and long pants, socks and shoes, plus gloves that are resistant to chlorine. If you have sensitive skin, wear a protective suit. If you have sensitive skin, wear a protective suit. If you have sensitive skin, wear a protective suit.

Engineering Control Statements:

When handling use closed systems, enclosed areas, or enclosures in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(a)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

Use Safety Recommendations

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets on skin. Then wash thoroughly and put on clean clothing.
- Remove and wash contaminated clothing before reuse.
- Rinse PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. Do not runoff from treated areas may be hazardous to fish and aquatic organisms.

In adjacent aquatic sites. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only trained handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

This product may be used in accordance with the labeling with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry sites. All the requirements in this label only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for entry into a treated area that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water: 1) clothing over long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves made of any synthetic material such as nitrile or rubber; 2) 14 mils.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this label only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when the product is used to produce agricultural plants for future, commercial purposes.

Do NOT enter or allow others to enter treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross contamination with other pesticides.

Pesticide Disposal: Pesticide wastes may be toxic. Improper disposal of unused pesticides, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance on proper disposal methods.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple-rinse container or equipment promptly after emptying. Triple-rinse as follows: Empty the container by tilting into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour into application equip-

STORAGE AND DISPOSAL (continued)

ment or a mix tank or show results for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. After triple rinsing procedure dispose of container.

GENERAL INFORMATION

Chlorpyrifos FLO is a broad spectrum fungicide with preventive, systemic, residual and curative properties recommended for the control or suppression of certain turf diseases. Disease control is optimal when it is applied in a preventive and scheduled spray program and used in rotation with other fungicides.

Failure to follow these label instructions may result in turfgrass injury and inferior turfgrass quality. This product may cause staining. Avoid off-target sites such as sidewalks, patios, driveways, lawns or similar surfaces. Avoid entry to areas used for recreation and sports until sprays have completely dried after application or prevent contact of clothing and shoes.

Chlorpyrifos FLO can be used on all commercial, institutional and residential turfgrass, such as golf courses, lawns, commercial lawns, rights of way, parks, recreation areas and sports fields, but for use by homeowners.

APPLICATION INFORMATION

Chlorpyrifos FLO may be used alone to control important diseases of turfgrass or alternatively used in tank mix or sequentially with other registered turfgrass fungicides.

Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control. Do not make applications when conditions favor drift beyond the target application area. Applications of Chlorpyrifos FLO must be made in accordance with the air drift controls of the label.

RESISTANCE MANAGEMENT

The active ingredient in Chlorpyrifos FLO belongs to the Sterol Inhibitors or Demethylase Inhibitors class of chemistry (Group 3). To maintain long term effectiveness of the fungicide, follow the specific resistance management guidance from a local extension specialist.

To limit the potential for development of disease resistance to the fungicide class do not make more than 3 sequential applications of Chlorpyrifos FLO or any Group 3 containing fungicide based rotating with a fungicide from a different group.

GENERAL RESTRICTIONS AND LIMITATIONS

DO NOT apply where the maximum annual rate from any combination of products containing chlorpyrifos are in the range 22.465 F. Repeat applications at 10-14 day intervals when heavy disease pressure is anticipated. For golf courses only, DO NOT apply to turf higher than 1 inch on golf holes when water bodies are present.

SPRAY MIXING AND COMPATIBILITY

In general, begin with clean application equipment and add one half of the required quantity of water to the spray mixing tank and start agitation. Refer to the product amounts tabulated under PRODUCT QUANTITY (FLO) for more information.

BY SPRAY MIXING AND TANK CAPACITY and add the required quantity of fungicide and the tank-mix carrier if applicable to the water and complete filling with water to the required total volume. Follow the recommendations of your State Cooperative Extension Service for tank mixing with other products. To prevent fuel-line or water-soluble granular products, liquid fertilizers and slow-release fertilizers and other herbicides from clogging the sprayer, flush the sprayer with water after every mixture run in the overnight or for long periods during the day without application.

The turf safety of general lawns mixes with Chlorpyrifos FLO including wettable and other products has not been tested on all turf types. Before applying any tank-mixes not specifically recommended on the label, safety to the target turf should be confirmed. Chlorpyrifos FLO is compatible with most commonly used fungicides, herbicides, insecticide, growth regulator and foliar nutrient products. However, the physical compatibility of Chlorpyrifos FLO with all general lawns mix partners has not been fully investigated. If tank mixing with other products is desirable, conduct a jar test with the volumes and rates typically used in agricultural application. Using a small container with water, thoroughly mix the products, wettable granules and slow-release granular products first. Then liquid fertilizers are immediately incorporated into. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, foam, or other procedures. Physical compatibility is assumed if the combination remains mixed or can be removed easily.

The use of soil penetrating adjuvant may improve the movement into the soil when irrigation after treatment is required for disease control.

Combination of Chlorpyrifos FLO at high labeled rates with PGRs may impact ball quality and reduce turf growth particularly during periods of heat stress and high humidity.

When preparing spray mixtures of Chlorpyrifos FLO with Super Products and the products in the following table, 1) avoid spraying products, 2) Chlorpyrifos FLO followed by other sprays, and 3) Chlorpyrifos FLO followed by other sprays.

RESISTANCE MANAGEMENT

The active ingredient in Chlorpyrifos FLO belongs to the Sterol Inhibitors or Demethylase Inhibitors class of chemistry (Group 3). To maintain long term effectiveness of the fungicide, follow the specific resistance management guidance from a local extension specialist.

To limit the potential for development of disease resistance to the fungicide class do not make more than 3 sequential applications of Chlorpyrifos FLO or any Group 3 containing fungicide based rotating with a fungicide from a different group.

GENERAL RESTRICTIONS AND LIMITATIONS

DO NOT apply where the maximum annual rate from any combination of products containing chlorpyrifos are in the range 22.465 F. Repeat applications at 10-14 day intervals when heavy disease pressure is anticipated. For golf courses only, DO NOT apply to turf higher than 1 inch on golf holes when water bodies are present.

SPRAY MIXING AND COMPATIBILITY

In general, begin with clean application equipment and add one half of the required quantity of water to the spray mixing tank and start agitation. Refer to the product amounts tabulated under PRODUCT QUANTITY (FLO) for more information.

TURF-SPECIFIC USE DIRECTIONS*

Chlorpyrifos FLO controls the diseases listed below.

Disease Control	Rate (lb ai Product / 1,000 sq ft)	Use Information
Anthracnose (Colletotrichum cereale)	0.4 - 1.1	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed on a 14-28 day interval. Under conditions for severe disease or for an early curative application, use the higher rate and shorter interval.
Brown Patch (Rhizoctonia solani)	0.4 - 1.1	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed on a 14-28 day interval. Under conditions for severe disease or for an early curative application, use the higher rate and shorter interval.
Brown Patch, Cool weather/Yellow patch (Rhizoctonia cereale)	0.5 - 1.1	Make 1-2 applications in the fall using a 21-28 day interval or when conditions are favorable for disease development.
Brown Ring (Wheat) Patch (Blumeria circinata var. cynodoni)	0.5 - 1.1	Begin fungicide applications at the early stage of yellow ring development when conditions are favorable for disease development. Continue as needed on a 14-28 day interval. Under conditions for severe disease or for an early curative application, use the higher rate and shorter interval. Rotate with 25 FIF fungicide in suggested for resistance management.
Dollar Spot (Sclerotinia homocarpa)	0.28 - 1.1	Begin fungicide applications preventively. Continue as needed on a 14-28 day interval. Under preventive applications when conditions are favorable for disease development, continue as needed on a 14-28 day interval. Under conditions for severe disease or for an early curative application, use the higher rate and shorter interval. Rotate with 25 FIF fungicide in suggested for resistance management.
Maximum Patch (Microstroma healei)	0.28 - 1.1	Begin fungicide applications preventively in the spring and early summer when the turf is cool and temperatures are in the range 22-65 F. Repeat applications at 10-14 day intervals when heavy disease pressure is anticipated. Under conditions for severe disease use the higher rate and shorter interval.
Gray Snow Mold (Typharia blight) (Typharia spp.)	0.28 - 1.1	Begin fungicide applications preventively in the late fall. Make 1-2 applications, repeating at 14-28 day intervals when heavy disease pressure is anticipated. Under conditions for severe disease use the higher rate and shorter interval. Rotate with alternative modes of action that are effective on snow molds.

Continued

5.5"

5"

chipco TRITON® FLO

For control of diseases on turfgrasses of commercial, institutional and residential turfgrass, such as golf courses, sod farms, commercial lawns, rights of way, parks, recreation areas and sports fields.

ACTIVE INGREDIENT:

Triticonazole.....	30.1%
Other Ingredients.....	69.9%
Total	100.0%

* Contains 3 lbs triticonazole per US gallon
EPA Reg. No. 432-1487

KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY**
Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-800-331-2867

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Treat Symptomatically.

Net Contents
1 Gallon
79458762
79684088 090306A

**BACKED
by BAYER™**

GROUP 3 FUNGICIDE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross contamination with other pesticides.

Pesticide Disposal: Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. After triple rinsing procedure dispose of container.

Refer to attached leaflet for complete Directions for Use and Precautionary Statements.



Produced for:
Bayer Environmental Science
A Business Group of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

GROUP 3 FUNGICIDE

PULL HERE TO OPEN ►



chipco TRITON® FLO

For control of diseases on turfgrasses of commercial, institutional and residential turfgrass, such as golf courses, sod farms, commercial lawns, rights of way, parks, recreation areas and sports fields.

ACTIVE INGREDIENT:

Triticonazole..... 30.1%

OTHER INGREDIENTS 69.9%

Total 100.0%

* Contains 3 lbs triticonazole per US gallon

EPA Reg. No. 432-1487

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

For MEDICAL and TRANSPORTATION Emergencies
ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

Net Contents
1 Gallon

79458762
79684088 090306A

BACKED
by **BAYER™**

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Have person sip a glass of water if able to swallow.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
NOTE TO PHYSICIAN: Treat Symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear long sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves made of any waterproof material such as natural rubber \geq 14 mils (if more options are needed, follow the instructions for category A on an EPA chemical-resistance category selection chart).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove and wash contaminated clothing before reuse.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms

in adjacent aquatic sites. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves made of any of any waterproof material such as natural rubber \geq 14 mils.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when the product is used to produce agricultural plants on farms, nurseries or greenhouses.

DO NOT enter or allow others to enter treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross contamination with other pesticides.

Pesticide Disposal: Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equip-

STORAGE AND DISPOSAL (continued)

ment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. After triple rinsing procedure dispose of container.

GENERAL INFORMATION

Chipco TRITON® FLO is a broad spectrum fungicide with preventive, systemic, residual and curative properties recommended for the control or suppression of certain turf diseases. Disease control is optimal when it is applied in a preventive and scheduled spray program and used in rotation with other fungicides.

Failure to follow these label instructions may result in turfgrass injury and inferior turfgrass quality. This product may cause staining. Avoid off-target sites such as sidewalks, patios, driveways pavers or similar materials. Avoid entry to areas used for recreation and sports until spray deposits are completely dry after application to prevent staining of clothing and shoes.

Chipco TRITON FLO can be used on all commercial, institutional and residential turfgrass, such as golf courses, sod farms, commercial lawns, rights of way, parks, recreation areas and sports fields. Not for use by homeowners.

APPLICATION INFORMATION

Chipco TRITON FLO may be used alone to control important diseases of turfgrass or alternatively used in tank mix or sequentially with other registered turfgrass fungicides.

Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control. Do not make applications when conditions favor drift beyond the target application area. All applications of Chipco TRITON FLO must be made in accordance with the use directions of this label.

Chipco TRITON FLO, when used in conjunction with best turfgrass management practice, will control the diseases listed under **TURF-SPECIFIC USE DIRECTIONS**.

Chipco TRITON FLO, when combined with other products such as Chipco® Signature™, will relieve the symptoms of summer stress syndrome/decline.

Turfgrass Species:

Colonial and Creeping Bentgrass	Centipedegrass	Annual and Perennial Ryegrass
Common and Hybrid Bermudagrass	Dichondra	St. Augustinegrass
Annual, Rough and Kentucky Bluegrass	Fine and Tall Fescue	Zoysiagrass
Buffalograss	Seashore Paspalum	Kikuyugrass

DO NOT use on ultra-dwarf bermudagrass varieties.

AVOID the shorter application intervals if the turf is under excessive stress in the summer or under an intensive growth control program.

Ground Application

For ground application equipment use the appropriate product quantities described in this label in 1 to 5 gallons of water per 1,000 sq ft for turf (44 to 220 gallons per acre).

SPRAY MIXING AND COMPATIBILITY

In general, begin with clean spray equipment and add one-half of the required quantity of water to the spray or mixing tank and start agitation. Refer to the product amounts tabulated under **PRODUCT QUANTITY (FL OZ)**

by **SPRAY VOLUME** and **TANK CAPACITY** and add the required quantity of fungicide and the tank-mix partner if applicable to the water and complete filling with water to the required total volume. Follow the recommendations of your State Cooperative Extension Service for tank-mixing with other products. In general, follow the order beginning first with water soluble packaging (wait for it to completely dissolve), wettable powders and water-dispersible granular products, liquid flowables and suspension concentrates and emulsifiable concentrates last. Maintain agitation throughout spraying. Do not allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

The turf safety of all potential tank-mixes with Chipco TRITON FLO including additives and other pesticides has not been tested on all turf type variants. Before applying any tank-mixture not specifically recommended on this label, safety to the target turf should be confirmed. Chipco TRITON FLO is compatible with most commonly used fungicide, herbicide, insecticide, growth regulator and foliar nutrient products. However, the physical compatibility of Chipco TRITON FLO with all potential tank-mix partners has not been fully investigated. If tank mixing with other pesticides is desirable, conduct a jar test with the volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products: wettable powders and water-dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

- The use of a soil penetrating adjuvant may improve the movement into the soil when irrigation after treatment is required for disease control.
- Combinations of Chipco TRITON FLO at high labeled rates with PGRs may impact turf quality and reduce turf growth particularly during period of heat stress and high humidity.
- When preparing spray mixtures of Chipco TRITON FLO with Bayer Products add the products in the following order: 1) Water dispersible products, 2) Chipco TRITON FLO followed by other suspension concentrates such as Compass®, 26 GT®, and lastly 3) Chipco® Signature™.

RESISTANCE MANAGEMENT

The active ingredient in Chipco TRITON FLO belongs to the Sterol Inhibitors or Demethylation Inhibitors class of chemistry (Group 3). To maintain long term effectiveness of this fungicide, follow the specific resistance management guidance from a local extension specialist.

To limit the potential for development of disease resistance to this fungicide class do not make more than 3 sequential applications of Chipco TRITON FLO or any Group 3 containing fungicide before rotating with a fungicide from a different group.

GENERAL RESTRICTIONS AND LIMITATIONS

DO NOT apply more than the maximum seasonal rate from any combination of products containing triticonazole.

DO NOT apply more than 3.3 fl oz of Chipco TRITON FLO per 1,000 sq ft (3.451 lb ai/A) per year.

Follow the **TURF-SPECIFIC USE DIRECTIONS** for repeat application intervals.

For golf courses only: DO NOT apply to turf cut higher than 1 inch on golf holes where water bodies are present.

This product cannot be used to formulate or reformulate any other pesticide product.

DO NOT apply this product to crops other than turfgrass.

DO NOT apply to areas used for grazing.

DO NOT feed clippings to livestock or poultry.

DO NOT apply through irrigation equipment of any type.

DO NOT use on ultra-dwarf bermudagrass varieties.

DO NOT use on ultradwarf bermudagrass varieties.

TURF-SPECIFIC USE DIRECTIONS¹

Chipco TRITON FLO controls the diseases listed below.

Disease Control	Rate fl oz Product /1,000 sq ft	Use Information
Anthracnose (<i>Colletotrichum cereale</i>)	0.41 - 1.1	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. Under conditions for severe disease or for an early curative application , use the higher rate and shorter interval.
Brown Patch (<i>Rhizoctonia solani</i>)	0.41 - 1.1	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. Under conditions for severe disease or for an early curative application , use the higher rate and shorter interval.
Brown Patch, Cool weather/Yellow patch (<i>Rhizoctonia cerealis</i>)	0.55 - 1.1	Make 1 to 2 applications in the fall using a 21-28 day interval or when conditions are favorable for disease development.
Brown Ring (Waitea) Patch (<i>Waitea circinata</i> var. <i>circinata</i>)	0.55 - 1.1	Begin fungicide applications at the early stage of yellow ring development when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. Late curative applications will not be effective. Irrigate to move the fungicide beneath the thatch.
Dollar Spot (<i>Sclerotinia homeocarpa</i>)	0.28 - 1.1	Begin fungicide applications preventively. Continue as needed, on a 14-28 day interval. Under preventive applications where light disease pressure is anticipated, use the lower rate and longer interval. Under conditions for severe disease or for an early curative application , use the higher rate and shorter interval. Rotation with 26 GT [®] Fungicide is suggested for resistance management.
Fusarium Patch (<i>Microdochium nivale</i>)	0.28 - 1.1	Begin fungicide applications preventively in the spring and early summer when the turf stays moist and temperatures are in the range 32-65° F. Repeat applications at 10-14 day intervals when heavy disease pressure is anticipated. Under conditions for severe disease use the higher rate and tank mix with 26 GT Fungicide.
Gray Snow Mold, Typhula Blight (<i>Typhula</i> spp.)	0.28 - 1.1	Begin fungicide applications preventively in the late fall. Make 1-2 applications, repeating at 14-28 day intervals when heavy disease pressure is anticipated. Under conditions for severe disease use the higher rate and tank mix with other fungicides with alternative modes of action that are effective on snow molds.

Chipco TRITON FLO at 1.1 fluid ounces per 1,000 sq ft

SPRAY VOLUME (gallons per 1,000 sq ft)

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons	
Spray Tank Capacity	25 Gallons	28 fl oz	14 fl oz	9.3 fl oz	7 fl oz	5.6 fl oz
	50 Gallons	56 fl oz	28 fl oz	18.7 fl oz	14 fl oz	11.2 fl oz
	100 Gallons	112 fl oz	56 fl oz	37.3 fl oz	28 fl oz	22.4 fl oz
	200 Gallons	224 fl oz	112 fl oz	74.6 fl oz	56 fl oz	44.8 fl oz

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), Chipco®, Chipco Triton®, Compass®, 26 GT®, Signature™, and Backed by Bayer™ are trademarks of Bayer.



Produced for:

Bayer Environmental Science

A Business Group of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709